


WHAT IS CLAIMED IS:

1. A compound of Formula (I):



or a stereoisomer or a pharmaceutically acceptable salt form thereof, wherein:

R^1 is selected from

- 10 H, $C(=O)R^{2a}$, $C(=O)OR^{2a}$, $S(=O)R^{2a}$, $S(=O)_2R^{2a}$,
C₃₋₇ cycloalkyl,
C₁₋₄ alkyl substituted with 0-3 R^2 ,
C₂₋₄ alkenyl substituted with 0-2 R^2 ,
C₂₋₄ alkynyl substituted with 0-2 R^2 ,
- 15 aryl substituted with 0-5 R^{42} ,
C₃₋₁₀ carbocyclic residue substituted with 0-3 R^{41} , and
5-6 membered heterocyclic ring system containing from 1-4 heteroatoms
selected from the group consisting of N, O, and S substituted with 0-3
 R^{41} ;

20

R^2 , at each occurrence, is independently selected from

- halo, C₁₋₃ haloalkyl, C₁₋₄ alkoxy, C₁₋₄ alkyl,
C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₃₋₆ cycloalkyl,
aryl substituted with 0-5 R^{42} ;
- 25 C₃₋₁₀ carbocyclic residue substituted with 0-3 R^{41} , and

5-6 membered heterocyclic ring system containing from 1-4 heteroatoms
selected from the group consisting of N, O, and S substituted with 0-3
R⁴¹;

- 5 R^{2a} is H, C₁₋₄ alkyl, (aryl)C₁₋₄ alkyl-, or
(C₃₋₆ cycloalkyl)C₁₋₄ alkyl-;

R^{4a} is H or C₁₋₄ alkyl;

- 10 R⁵ is H, C₁₋₄ alkyl substituted with 0-2 R²⁰,
-C(=O)(C₁₋₄ alkyl), -C(=O)O(C₁₋₄ alkyl), or C₁₋₄ haloalkyl;

R⁶ is selected from

- 15 halo, -CF₃, -OCF₃, -CN, -NO₂, -OCH₃, -SCH₃, -CF₂CF₃, -O-R¹¹,
-OCF₂CF₃, -OCF₂H, -OCF₂CH₃,
-S-R¹¹, -S(=O)-R¹¹, -S(=O)₂-R¹¹, -S(=O)-NR¹⁰-R¹¹,
-S(=O)₂-NR¹⁰-R¹¹, -NR¹⁰-R¹¹, -CH₂O-R¹¹, -CH₂S-R¹¹,
CH₂S(=O)-R¹¹, CH₂S(=O)₂-R¹¹, -CH₂NR¹⁰-R¹¹, -C(=O)NR¹⁰-R¹¹
C₁₋₄ haloalkyl, (C₁₋₄ haloalkyl)oxy;
20 C₁₋₄ alkyl substituted with 0-2 R²⁰,
C₂₋₄ alkenyl substituted with 0-2 R²⁰,
C₂₋₄ alkynyl substituted with 0-1 R²⁰, and
C₃₋₆ carbocyclic residue substituted with 0-3 R²¹,

- 25 R⁷ and R⁹ are independently selected from
H, F, Cl, Br, -CF₃, -OCF₃, -OH, -CN, -NO₂, -CF₂CF₃, C₁₋₄ alkyl,
C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and
(C₁₋₄ haloalkyl)oxy;

R⁸ is selected from

- halo, -CF₃, -OCF₃, -OH, -CN, -NO₂, -OCH₃, -SCH₃, -CF₂CF₃,
 -OR¹², -SR¹², -NR¹²R¹³, -C(O)H, -C(O)R¹², -C(O)NR¹²R¹³,
 -NR¹⁴C(O)R¹², -C(O)OR¹², -OC(O)R¹², -OC(O)OR¹²,
 5 -S(O)R¹², -S(O)₂R¹², -S(O)NR¹²R¹³, -S(O)₂NR¹²R¹³,
 -NR¹⁴S(O)R¹², -NR¹⁴S(O)₂R¹², -NR¹²C(O)R¹⁵, -NR¹²C(O)OR¹⁵,
 -NR¹²S(O)₂R¹⁵, -NR¹²C(O)NHR¹⁵;
 C₁₋₆ alkyl substituted with 0-2 R^{8a},
 C₂₋₆ alkenyl substituted with 0-2 R^{8a},
 10 C₂₋₆ alkynyl substituted with 0-2 R^{8a},
 C₃₋₆ cycloalkyl substituted with 0-2 R^{8a},
 C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³;

R^{8a}, at each occurrence, is independently selected from

- 15 halo, -CF₃, -OCF₃, -OH, -CN, -NO₂, -CF₂CF₃,
 methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl,
 -OR¹², -SR¹², -NR¹²R¹³, -C(O)H, -C(O)R¹², -C(O)NR¹²R¹³,
 -NR¹⁴C(O)R¹², -C(O)OR¹², -OC(O)R¹², -OC(O)OR¹²,
 -S(O)R¹², -S(O)₂R¹², -S(O)NR¹²R¹³, -S(O)₂NR¹²R¹³,
 20 -NR¹⁴S(O)R¹², -NR¹⁴S(O)₂R¹², -NR¹²C(O)R¹⁵, -NR¹²C(O)OR¹⁵,
 -NR¹²S(O)₂R¹⁵, -NR¹²C(O)NHR¹⁵;
 phenyl substituted with 0-5 R³³;
 C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³, and
 5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
 25 selected from the group consisting of N, O, and S substituted with 0-3
 R³³;

R¹⁰ is H or C₁₋₄ alkyl;

R¹¹ is selected from

- C₁₋₆ alkyl substituted with 0-2 R²⁰,
- C₂₋₆ alkenyl substituted with 0-2 R²⁰,
- 5 C₂₋₆ alkynyl substituted with 0-1 R²⁰,
- C₃₋₁₀ carbocyclic residue substituted with 0-3 R²¹,
- aryl substituted with 0-5 R²³, and
- 5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
selected from the group consisting of N, O, and S substituted with 0-3
10 R²¹;

alternatively, R¹⁰ and R¹¹ join to form a 5- or 6-membered ring optionally
substituted with -O- or -N(R¹⁴)-;

- 15 alternatively, R¹⁰ and R¹¹ when attached to N may be combined to form a 9- or 10-
membered bicyclic heterocyclic ring system containing from 1-3 heteroatoms
selected from the group consisting of N, O, and S, wherein said bicyclic
heterocyclic ring system is unsaturated or partially saturated, wherein said
bicyclic heterocyclic ring system is substituted with 0-3 R¹⁶;

20

R¹² is selected from H,

- C₁₋₆ alkyl substituted with 0-2 R^{12a},
- C₂₋₆ alkenyl substituted with 0-2 R^{12a},
- C₂₋₆ alkynyl substituted with 0-2 R^{12a},
- 25 C₃₋₆ cycloalkyl substituted with 0-3 R³³,
- aryl substituted with 0-5 R³³;
- C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³, and

5-10 membered heterocyclic ring system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-3 R³³;

5 R^{12a}, at each occurrence, is independently selected from
 H, halo, -OH, -CN, -NO₂, -CO₂H, -SO₂R⁴⁵, -SOR⁴⁵, -SR⁴⁵,
 -NR⁴⁶SO₂R⁴⁵, -NR⁴⁶COR⁴⁵, -NR⁴⁶R⁴⁷, -SO₂NR⁴⁶R⁴⁷,
 -CONR⁴⁶R⁴⁷, -OR⁴⁵, =O,
 C₁₋₄ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl,
 10 phenyl substituted with 0-5 R³³;
 C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³, and
 5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
 selected from the group consisting of N, O, and S substituted with 0-3
 R³³;

15 R¹³, at each occurrence, is independently selected from
 H, C₁₋₄ alkyl, C₂₋₄ alkenyl, and C₂₋₄ alkynyl;

alternatively, R¹² and R¹³ join to form a 5- or 6-membered ring optionally
 20 substituted with -O- or -N(R¹⁴)-;

alternatively, R¹² and R¹³ when attached to N may be combined to form a 9- or 10-
 membered bicyclic heterocyclic ring system containing from 1-3 heteroatoms
 selected from the group consisting of N, O, and S, wherein said bicyclic
 25 heterocyclic ring system is unsaturated or partially saturated, wherein said
 bicyclic heterocyclic ring system is substituted with 0-3 R¹⁶;

R¹⁴, at each occurrence, is independently selected from H and C₁₋₄ alkyl;

30 R¹⁵, at each occurrence, is independently selected from

H, C₁₋₄ alkyl, C₂₋₄ alkenyl, and C₂₋₄ alkynyl;

R¹⁶, at each occurrence, is independently selected from

H, OH, halo, CN, NO₂, CF₃, SO₂R⁴⁵, NR⁴⁶R⁴⁷, -C(=O)H,
 5 C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ haloalkyl,
 C₁₋₃ haloalkyl-oxy-, and C₁₋₃ alkyloxy-;

R²⁰ is selected from

H, halo, -OH, -CF₃, -CN, -NO₂, -CO₂H, -SO₂R⁴⁵,
 10 -SOR⁴⁵, -SR⁴⁵, -NR⁴⁶SO₂R⁴⁵, -NR⁴⁶COR⁴⁵, -NR⁴⁶R⁴⁷,
 C₁₋₄ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₄ alkoxy,
 C₁₋₄ haloalkyl;
 C₃₋₁₀ carbocyclic residue substituted with 0-3 R²¹;
 aryl substituted with 0-5 R²³; and
 15 5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
 selected from the group consisting of N, O, and S substituted with 0-3
 R²¹;

R²¹, at each occurrence, is independently selected from

20 H, OH, halo, CF₃, SO₂R⁴⁵, NR⁴⁶R⁴⁷, CN, NO₂, =O, C₁₋₄ alkyl,
 C₁₋₄ alkoxy, and (C₁₋₄ haloalkyl)oxy;

R²³, at each occurrence, is independently selected from

H, OH, halo, CF₃, SO₂R⁴⁵, NR⁴⁶R⁴⁷, CN, NO₂, C₁₋₄ alkyl,
 25 C₁₋₄ alkoxy, and (C₁₋₄ haloalkyl)oxy;

R³³, at each occurrence, is independently selected from

H, OH, halo, -CN, -NO₂, -CF₃, -OCF₃, -SO₂R³⁵, -S(=O)R³⁵, -SR³⁵,
 -NR³⁶R³⁷, -NHC(=O)R³⁵, -C(=O)NR³⁶R³⁷, -C(=O)H, -C(=O)R³⁵,

-C(=O)OR³⁵, -OC(=O)R³⁵, -OR³⁵,
 C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₄ haloalkyl,
 C₁₋₄ alkoxy, (C₁₋₄ haloalkyl)oxy,
 C₃₋₆ cycloalkyl, phenyl, aryl substituted with 0-2 R³⁴,
 5 C₁₋₆ alkyl substituted with R³⁴, and
 C₂₋₆ alkenyl substituted with R³⁴;

R³⁴, at each occurrence, is independently selected from
 OH, C₁₋₄ alkoxy, -SO₂R³⁵, -NR³⁶R³⁷, NR³⁶R³⁷C(=O)-, and
 10 (C₁₋₄ alkyl)CO₂-;

R³⁵, at each occurrence, is independently selected from
 C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl,
 (C₃₋₆ cycloalkyl)methyl-, and (C₃₋₆ cycloalkyl)ethyl-;

15 R³⁶, at each occurrence, is independently selected from H and C₁₋₄ alkyl;

R³⁷, at each occurrence, is independently selected from H, C₁₋₄ alkyl,
 -C(=O)NH(C₁₋₄ alkyl), -SO₂(C₁₋₄ alkyl),
 20 -C(=O)O(C₁₋₄ alkyl), -C(=O)(C₁₋₄ alkyl), and -C(=O)H;

R⁴¹, at each occurrence, is independently selected from
 H, CF₃, halo, OH, CO₂H, SO₂R⁴⁵, NR⁴⁶R⁴⁷, NO₂, CN, =O,
 C₁₋₄ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkyl;

25 R⁴², at each occurrence, is independently selected from
 H, CF₃, halo, OH, CO₂H, SO₂R⁴⁵, SOR⁴⁵, SR⁴⁵, NR⁴⁶SO₂R⁴⁵,
 NR⁴⁶COR⁴⁵, NR⁴⁶R⁴⁷, NO₂, CN,
 C₁₋₄ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkyl;

R⁴⁵ is C₁₋₄ alkyl;

R⁴⁶, at each occurrence, is independently selected from H and C₁₋₄ alkyl;

5

R⁴⁷, at each occurrence, is independently selected from H, C₁₋₄ alkyl,

-C(=O)NH(C₁₋₄ alkyl), -SO₂(C₁₋₄ alkyl),

-C(=O)O(C₁₋₄ alkyl), -C(=O)(C₁₋₄ alkyl), and -C(=O)H;

10 m is 1 or 2;


provided that when R¹¹ is C₁₋₆ alkyl, then R¹ is not a C₁₋₄ alkyl substituted by a) an unsubstituted 3H-pyrimidine-4-one moiety, b) a substituted 3H-pyrimidine-4-one moiety, c) an unsubstituted bicyclic derivative of 3H-pyrimidine-4-one, or d) a substituted bicyclic derivative of 3H-pyrimidine-4-one;

15

provided that when R⁶ is -O-R¹¹ and R⁶ is C₁₋₆ alkyl; then R^{8a} is not a substituted or unsubstituted indole moiety.

20

2. A compound of Claim 1 of Formula (Ia):



25 or a stereoisomer or a pharmaceutically acceptable salt form thereof, wherein:

R¹ is selected from

H, C₁₋₃ haloalkyl, C₃₋₆ cycloalkyl,
 C₁₋₄ alkyl substituted with 0-2 R²,
 C₂₋₄ alkenyl substituted with 0-2 R², and
 5 C₂₋₄ alkynyl substituted with 0-2 R²;

R², at each occurrence, is independently selected from

halo, C₁₋₃ haloalkyl, C₁₋₄ alkoxy, C₁₋₄ alkyl,
 C₃₋₆ cycloalkyl, and phenyl substituted with 0-5 R⁴²;

10

R^{4a} is H or C₁₋₄ alkyl;

R⁵ is H, C₁₋₄ alkyl substituted with 0-1 R²⁰, or C₁₋₄ haloalkyl;

15 R⁶ is selected from

halo, -CF₃, -OCF₃, -CN, -NO₂, -OCH₃, -SCH₃, -CF₂CF₃, -O-R¹¹,
 -OCF₂CF₃, -OCF₂H, -OCF₂CH₃,
 -S-R¹¹, -S(=O)-R¹¹, -S(=O)₂-R¹¹, -NR¹⁰-R¹¹, -CH₂O-R¹¹,
 -CH₂S-R¹¹, CH₂S(=O)-R¹¹, CH₂S(=O)₂-R¹¹, -CH₂NR¹⁰-R¹¹,
 20 C₁₋₄ haloalkyl, (C₁₋₄ haloalkyl)oxy;
 C₁₋₄ alkyl substituted with 0-2 R²⁰,
 C₂₋₄ alkenyl substituted with 0-2 R²⁰,
 C₂₋₄ alkynyl substituted with 0-1 R²⁰, and
 C₃₋₆ carbocyclic residue substituted with 0-3 R²¹,

25

R⁷ and R⁹ are independently selected from

H, F, Cl, Br, -CF₃, -OCF₃, -OH, -CN, -NO₂, CF₂CF₃, C₁₋₄ alkyl,
 C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and

(C₁₋₄ haloalkyl)oxy;

R⁸ is selected from

- halo, -CF₃, -OCF₃, -OH, -CN, -NO₂, -OCH₃, -SCH₃, -CF₂CF₃,
 5 -OR¹², -SR¹², -NR¹²R¹³, -C(O)H, -C(O)R¹², -C(O)NR¹²R¹³,
 -NR¹⁴C(O)R¹², -C(O)OR¹², -OC(O)R¹², -OC(O)OR¹²,
 -S(O)R¹², -S(O)₂R¹², -S(O)NR¹²R¹³, -S(O)₂NR¹²R¹³,
 -NR¹⁴S(O)R¹², -NR¹⁴S(O)₂R¹², -NR¹²C(O)R¹⁵, -NR¹²C(O)OR¹⁵,
 -NR¹²S(O)₂R¹⁵, -NR¹²C(O)NHR¹⁵;
 10 C₁₋₆ alkyl substituted with 0-2 R^{8a},
 C₂₋₆ alkenyl substituted with 0-2 R^{8a},
 C₂₋₆ alkynyl substituted with 0-2 R^{8a},
 C₃₋₆ cycloalkyl substituted with 0-2 R^{8a}, and
 C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³;

15

R^{8a}, at each occurrence, is independently selected from

- halo, -CF₃, -OCF₃, -OH, -CN, -NO₂, -CF₂CF₃,
 methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl,
 -OR¹², -SR¹², -NR¹²R¹³, -C(O)H, -C(O)R¹², -C(O)NR¹²R¹³,
 20 -NR¹⁴C(O)R¹², -C(O)OR¹², -OC(O)R¹², -OC(O)OR¹²,
 -S(O)R¹², -S(O)₂R¹², -S(O)NR¹²R¹³, -S(O)₂NR¹²R¹³,
 -NR¹⁴S(O)R¹², -NR¹⁴S(O)₂R¹², -NR¹²C(O)R¹⁵, -NR¹²C(O)OR¹⁵,
 -NR¹²S(O)₂R¹⁵, -NR¹²C(O)NHR¹⁵;

phenyl substituted with 0-5 R³³;

25

C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³, and

5-6 membered heterocyclic ring system containing from 1-4 heteroatoms

selected from the group consisting of N, O, and S substituted with 0-3

R³³;

R¹⁰ is H or C₁₋₄ alkyl;

R¹¹ is selected from

- 5 C₁₋₆ alkyl substituted with 0-2 R²⁰,
 C₂₋₆ alkenyl substituted with 0-2 R²⁰,
 C₂₋₆ alkynyl substituted with 0-1 R²⁰,
 C₃₋₁₀ carbocyclic residue substituted with 0-3 R²¹,
 aryl substituted with 0-5 R²³, and
10 5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
 selected from the group consisting of N, O, and S substituted with 0-3
 R²¹;

- alternatively, R¹⁰ and R¹¹ join to form a 5- or 6-membered ring optionally
15 substituted with -O- or -N(R¹⁴)-;

- alternatively, R¹⁰ and R¹¹ when attached to N may be combined to form a 9- or 10-
 membered bicyclic heterocyclic ring system containing from 1-3 heteroatoms
 selected from the group consisting of N, O, and S, wherein said bicyclic
20 heterocyclic ring system is unsaturated or partially saturated, wherein said
 bicyclic heterocyclic ring system is substituted with 0-3 R¹⁶;

R¹² is selected from H,

- C₁₋₆ alkyl substituted with 0-2 R^{12a},
25 C₂₋₆ alkenyl substituted with 0-2 R^{12a},
 C₂₋₆ alkynyl substituted with 0-2 R^{12a},
 C₃₋₆ cycloalkyl substituted with 0-3 R³³,
 aryl substituted with 0-5 R³³;
 C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³, and

5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
selected from the group consisting of N, O, and S substituted with 0-3
R³³;

5 R^{12a}, at each occurrence, is independently selected from
H, halo, -OH, -CN, -NO₂, -CO₂H, -SO₂R⁴⁵, -SOR⁴⁵, -SR⁴⁵,
-NR⁴⁶SO₂R⁴⁵, -NR⁴⁶COR⁴⁵, -NR⁴⁶R⁴⁷, -SO₂NR⁴⁶R⁴⁷,
-CONR⁴⁶R⁴⁷, -OR⁴⁵, =O,
C₁₋₄ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl,
10 phenyl substituted with 0-5 R³³;
C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³, and
5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
selected from the group consisting of N, O, and S substituted with 0-3
R³³;

15 R¹³, at each occurrence, is independently selected from
H, C₁₋₄ alkyl, C₂₋₄ alkenyl, and C₂₋₄ alkynyl;

alternatively, R¹² and R¹³ join to form a 5- or 6-membered ring optionally
20 substituted with -O- or -N(R¹⁴)-;

alternatively, R¹² and R¹³ when attached to N may be combined to form a 9- or 10-
membered bicyclic heterocyclic ring system containing from 1-3 heteroatoms
selected from the group consisting of N, O, and S, wherein said bicyclic
25 heterocyclic ring system is unsaturated or partially saturated, wherein said
bicyclic heterocyclic ring system is substituted with 0-3 R¹⁶;

R¹⁴, at each occurrence, is independently selected from H and C₁₋₄ alkyl;

30 R¹⁵, at each occurrence, is independently selected from

H, C₁₋₄ alkyl, C₂₋₄ alkenyl, and C₂₋₄ alkynyl;

R¹⁶, at each occurrence, is independently selected from

H, OH, halo, CN, NO₂, CF₃, SO₂R⁴⁵, NR⁴⁶R⁴⁷, -C(=O)H,
 5 C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ haloalkyl,
 C₁₋₃ haloalkyl-oxy-, and C₁₋₃ alkyloxy-;

R²⁰ is selected from

H, halo, -OH, -CF₃, -CN, -NO₂, -CO₂H, -SO₂R⁴⁵,
 10 -SOR⁴⁵, -SR⁴⁵, -NR⁴⁶SO₂R⁴⁵, -NR⁴⁶COR⁴⁵, -NR⁴⁶R⁴⁷,
 C₁₋₄ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₄ alkoxy,
 C₁₋₄ haloalkyl;
 C₃₋₁₀ carbocyclic residue substituted with 0-3 R²¹;
 aryl substituted with 0-5 R²³; and
 15 5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
 selected from the group consisting of N, O, and S substituted with 0-3
 R²¹;

R²¹, at each occurrence, is independently selected from

20 H, OH, halo, CF₃, SO₂R⁴⁵, NR⁴⁶R⁴⁷, CN, NO₂, =O, C₁₋₄ alkyl;
 C₁₋₄ alkoxy, and (C₁₋₄ haloalkyl)oxy;

R²³, at each occurrence, is independently selected from

H, OH, halo, CF₃, SO₂R⁴⁵, NR⁴⁶R⁴⁷, CN, NO₂, C₁₋₄ alkyl;
 25 C₁₋₄ alkoxy, and (C₁₋₄ haloalkyl)oxy;

R³³, at each occurrence, is independently selected from

H, OH, halo, -CN, -NO₂, -CF₃, -OCF₃, -SO₂R³⁵, -S(=O)R³⁵,
 -SR³⁵, -NR³⁶R³⁷, -NHC(=O)R³⁵, -C(=O)NR³⁶R³⁷,

$-\text{C}(=\text{O})\text{H}$, $-\text{C}(=\text{O})\text{R}^{35}$, $-\text{C}(=\text{O})\text{OR}^{35}$, $-\text{OC}(=\text{O})\text{R}^{35}$, $-\text{OR}^{35}$,
 C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-4} haloalkyl,
 C_{1-4} alkoxy, $(\text{C}_{1-4} \text{ haloalkyl})\text{oxy}$,
 C_{3-6} cycloalkyl, phenyl, aryl substituted with 0-2 R^{34} ,
5 C_{1-6} alkyl substituted with R^{34} , and
 C_{2-6} alkenyl substituted with R^{34} ;

R^{34} , at each occurrence, is independently selected from
 OH , C_{1-4} alkoxy, $-\text{SO}_2\text{R}^{35}$, $-\text{NR}^{36}\text{R}^{37}$, $\text{NR}^{36}\text{R}^{37}\text{C}(=\text{O})-$, and
10 $(\text{C}_{1-4} \text{ alkyl})\text{CO}_2-$;

R^{35} , at each occurrence, is independently selected from
 C_{1-4} alkyl, C_{1-4} haloalkyl, C_{3-6} cycloalkyl,
 $(\text{C}_{3-6} \text{ cycloalkyl})\text{methyl-}$, and $(\text{C}_{3-6} \text{ cycloalkyl})\text{ethyl-}$;

15 R^{36} , at each occurrence, is independently selected from H and C_{1-4} alkyl;

R^{37} , at each occurrence, is independently selected from H, C_{1-4} alkyl,
 $-\text{C}(=\text{O})\text{NH}(\text{C}_{1-4} \text{ alkyl})$, $-\text{SO}_2(\text{C}_{1-4} \text{ alkyl})$,
20 $-\text{C}(=\text{O})\text{O}(\text{C}_{1-4} \text{ alkyl})$, $-\text{C}(=\text{O})(\text{C}_{1-4} \text{ alkyl})$, and $-\text{C}(=\text{O})\text{H}$;

R^{41} , at each occurrence, is independently selected from
H, CF_3 , halo, OH, CO_2H , SO_2R^{45} , $\text{NR}^{46}\text{R}^{47}$, NO_2 , CN, $=\text{O}$,
 C_{1-4} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{1-4} alkoxy, and C_{1-4} haloalkyl;

25 R^{42} , at each occurrence, is independently selected from
H, CF_3 , halo, OH, CO_2H , SO_2R^{45} , SOR^{45} , SR^{45} , $\text{NR}^{46}\text{SO}_2\text{R}^{45}$,
 $\text{NR}^{46}\text{COR}^{45}$, $\text{NR}^{46}\text{R}^{47}$, NO_2 , CN, C_{1-4} alkyl, C_{2-6} alkenyl,
 C_{2-6} alkynyl, C_{1-4} alkoxy, and C_{1-4} haloalkyl;

R⁴⁵ is C₁₋₄ alkyl;

R⁴⁶, at each occurrence, is independently selected from H and C₁₋₄ alkyl;

5


R⁴⁷, at each occurrence, is independently selected from H, C₁₋₄ alkyl,

-C(=O)NH(C₁₋₄ alkyl), -SO₂(C₁₋₄ alkyl),

-C(=O)O(C₁₋₄ alkyl), -C(=O)(C₁₋₄ alkyl), and -C(=O)H;

10 m is 1 or 2.

3. A compound of Claim 2 of Formula (Ia):



15

or a stereoisomer or a pharmaceutically acceptable salt form thereof, wherein:

R¹ is selected from

20

H, CF₃, methyl, ethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,

C₁₋₄ alkyl substituted with 0-1 R²,

C₂₋₄ alkenyl substituted with 0-1 R², and

C₂₋₄ alkynyl substituted with 0-1 R²;

25

R² is selected from

F, Cl, CH₂F, CHF₂, CF₃, methyl, ethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and phenyl;

R^{4a} is H or methyl;

5

R⁵ is H, methyl, or ethyl;

R⁶ is selected from

10 F, Cl, -CF₃, -OCF₃, -CF₂CF₃, -OCF₂CF₃, -OCF₂H, -OCF₂CH₃, -CN, -NO₂, -O-R¹¹, -S-R¹¹, -S(=O)-R¹¹, -S(=O)₂-R¹¹, -CH₂O-R¹¹, -CH₂S-R¹¹, CH₂S(=O)-R¹¹, CH₂S(=O)₂-R¹¹, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, and s-butyl;

R⁷ and R⁹ are independently selected from

15 H, F, Cl, -CH₃, -OCH₃, -CF₃, -OCF₃, -CN, and -NO₂;

R⁸ is selected from

-OR¹², -SR¹², -NR¹²R¹³, -C(O)R¹², -S(O)R¹², -S(O)₂R¹²,
C₁₋₆ alkyl substituted with 0-2 R^{8a},
20 C₂₋₆ alkenyl substituted with 0-2 R^{8a},
C₂₋₆ alkynyl substituted with 0-2 R^{8a},
C₃₋₆ cycloalkyl substituted with 0-2 R^{8a}, and
C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³;

25 R^{8a}, at each occurrence, is independently selected from

halo, -CF₃, -OCF₃, -OH, -CN, -NO₂, -CF₂CF₃,
methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl,
-OR¹², -SR¹², -NR¹²R¹³, -C(O)H, -C(O)R¹², -C(O)NR¹²R¹³,
-NR¹⁴C(O)R¹², -C(O)OR¹², -OC(O)R¹², -OC(O)OR¹²,

-S(O)R¹², -S(O)₂R¹², -S(O)NR¹²R¹³, -S(O)₂NR¹²R¹³,
 -NR¹⁴S(O)R¹², -NR¹⁴S(O)₂R¹², -NR¹²C(O)R¹⁵, -NR¹²C(O)OR¹⁵,
 -NR¹²S(O)₂R¹⁵, -NR¹²C(O)NHR¹⁵;

phenyl substituted with 0-5 R³³;

- 5 C₃-10 carbocyclic residue substituted with 0-3 R³³, and
 5-6 membered heterocyclic ring system containing from 1-4 heteroatoms
 selected from the group consisting of N, O, and S substituted with 0-3
 R³³;

- 10 R¹¹ is selected from
 methyl, ethyl, propyl, and phenyl substituted with 0-5 R²³,

R¹² is selected from

- C₁-6 alkyl substituted with 0-2 R^{12a},
 15 C₂-6 alkenyl substituted with 0-2 R^{12a},
 C₂-6 alkynyl substituted with 0-2 R^{12a},
 C₃-6 cycloalkyl substituted with 0-3 R³³,
 aryl substituted with 0-5 R³³;
 C₃-10 carbocyclic residue substituted with 0-3 R³³, and
 20 5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
 selected from the group consisting of N, O, and S substituted with 0-3
 R³³;

R^{12a}, at each occurrence, is independently selected from

- 25 H, halo, -OH, -CN, -NO₂, -CO₂H, -SO₂R⁴⁵, -SOR⁴⁵,
 -SR⁴⁵, -NR⁴⁶SO₂R⁴⁵, -NR⁴⁶COR⁴⁵, -NR⁴⁶R⁴⁷,
 -SO₂NR⁴⁶R⁴⁷, -CONR⁴⁶R⁴⁷, -OR⁴⁵, =O,
 C₁-4 alkyl, C₂-6 alkenyl, C₂-6 alkynyl,

phenyl substituted with 0-5 R³³;
C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³, and
5-10 membered heterocyclic ring system containing from 1-4 heteroatoms
selected from the group consisting of N, O, and S substituted with 0-3
5 R³³;

R¹³, at each occurrence, is independently selected from
H, C₁₋₄ alkyl, C₂₋₄ alkenyl, and C₂₋₄ alkynyl;

10 alternatively, R¹² and R¹³ join to form a 5- or 6-membered ring selected from
pyrrolyl, pyrrolidinyl, imidazolyl, piperidinyl, piperizinyl, methylpiperizinyl,
and morpholinyl;

alternatively, R¹² and R¹³ when attached to N may be combined to form a 9- or 10-
15 membered bicyclic heterocyclic ring system containing from 1-3 heteroatoms
selected from the group consisting of N, O, and S; wherein said bicyclic
heterocyclic ring system is selected from indolyl, indolinyl, indazolyl,
benzimidazolyl, benzimidazolinyl, and benztriazolyl; wherein said bicyclic
heterocyclic ring system is substituted with 0-1 R¹⁶;

20

R¹⁴ is H, methyl, ethyl, propyl, or butyl;

R¹⁵ is H, methyl, ethyl, propyl, or butyl;

25 R¹⁶, at each occurrence, is independently selected from
H, OH, F, Cl, CN, NO₂, methyl, ethyl, methoxy, ethoxy, trifluoromethyl, and
trifluoromethoxy;

R²³, at each occurrence, is independently selected from

H, OH, F, Cl, CF₃, SO₂R⁴⁵, NR⁴⁶R⁴⁷, CN, NO₂, methyl, ethyl, propyl, and butyl;

R³³, at each occurrence, is independently selected from

- 5 H, OH, halo, -CN, -NO₂, -CF₃, -OCF₃, -SO₂R³⁵, -S(=O)R³⁵,
 -SR³⁵, -NR³⁶R³⁷, -NHC(=O)R³⁵, -C(=O)NR³⁶R³⁷,
 -C(=O)H, -C(=O)R³⁵, -C(=O)OR³⁵, -OC(=O)R³⁵, -OR³⁵,
 C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₄ haloalkyl,
 C₁₋₄ alkoxy, (C₁₋₄ haloalkyl)oxy,
 10 C₃₋₆ cycloalkyl, phenyl, aryl substituted with 0-2 R³⁴,
 C₁₋₆ alkyl substituted with R³⁴, and
 C₂₋₆ alkenyl substituted with R³⁴;

R³⁴, at each occurrence, is independently selected from

- 15 OH, C₁₋₄ alkoxy, -SO₂R³⁵, -NR³⁶R³⁷, NR³⁶R³⁷C(=O)-, and (C₁₋₄
 alkyl)CO₂-;

R³⁵, at each occurrence, is independently selected from

- C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl,
 20 (C₃₋₆ cycloalkyl)methyl-, and (C₃₋₆ cycloalkyl)ethyl-;

R³⁶, at each occurrence, is independently selected from H and C₁₋₄ alkyl;

R³⁷, at each occurrence, is independently selected from H, C₁₋₄ alkyl,

- 25 -C(=O)NH(C₁₋₄ alkyl), -SO₂(C₁₋₄ alkyl),
 -C(=O)O(C₁₋₄ alkyl), -C(=O)(C₁₋₄ alkyl), and -C(=O)H;

R⁴⁵ is C₁₋₄ alkyl;

R⁴⁶, at each occurrence, is independently selected from H and C₁₋₄ alkyl;

R⁴⁷, at each occurrence, is independently selected from H, C₁₋₄ alkyl,


-C(=O)NH(C₁₋₄ alkyl), -SO₂(C₁₋₄ alkyl),

5 -C(=O)O(C₁₋₄ alkyl), -C(=O)(C₁₋₄ alkyl), and -C(=O)H;

m is 1 or 2.

4. A compound of Claim 3 of Formula (Ib):

10



or a stereoisomer or a pharmaceutically acceptable salt form thereof, wherein:

15

R¹ is selected from H, methyl, and ethyl;

R⁵ is H, methyl, or ethyl;

20 R⁶ is selected from

-F, -Cl, -CF₃, -OCF₃, -CF₂CF₃, -OCF₂CF₃, -OCF₂H, -OCF₂CH₃, -CN,
 -NO₂, -OCH₃, -OCH₂CH₃, -SCH₃, -SCH₂CH₃, -S(=O)CH₃,
 -S(=O)₂CH₃, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, and s-butyl;

25 R⁷ is H, F, or Cl;

R⁸ is selected from

- OR¹², -SR¹², -NR¹²R¹³, -C(O)R¹², -S(O)R¹², -S(O)₂R¹²,
- C₁₋₆ alkyl substituted with 0-2 R^{8a},
- C₃₋₆ cycloalkyl substituted with 0-2 R^{8a}, and
- 5 C₃₋₁₀ carbocyclic residue substituted with 0-3 R³³;

R^{8a}, at each occurrence, is independently selected from

- H, F, Cl, Br, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl,
- t-butyl, -OH, methoxy, ethoxy, n-propoxy, i-propoxy, -CF₃, -OCF₃,
- 10 -CN, -NO₂, -CF₂CF₃, -SCH₃, -SCH₂CH₃, -SO₂CH₃, -NH₂,
- CH₂NH(CH₃), -CH₂N(CH₃)₂, -NH(CH₃), -N(CH₃)₂, -CO(CH₃),
- CO(OCH₃), -NHCO(CH₃), -CONH₂, -C(=O)H, -CH(OH)CH₃, -CH₂OH,
- CH₂CH₂OH, -CH₂OCH₃, -CH₂CH₂OCH₃, -CH₂OCH₂CH₃,
- phenyl substituted with 0-5 R³³, and pyridyl substituted with 0-5 R³³

15

R⁹ is H;

R¹² is selected from

- C₁₋₆ alkyl substituted with 0-2 R^{12a},
- 20 cyclopropyl substituted with 0-2 R³³,
- cyclobutyl substituted with 0-2 R³³,
- cyclopentyl substituted with 0-2 R³³,
- cyclohexyl substituted with 0-2 R³³,
- bicyclo[3.1.1]heptane substituted with 0-2 R³³,
- 25 bicyclo[2.2.1]heptane substituted with 0-2 R³³,
- phenyl substituted with 0-3 R³³; and
- pyridyl substituted with 0-3 R³³;

R^{12a}, at each occurrence, is independently selected from


- H, F, Cl, -OH, methyl, ethyl,
 cyclopropyl substituted with 0-2 R³³,
 cyclobutyl substituted with 0-2 R³³,
 cyclopentyl substituted with 0-2 R³³,
 5 cyclohexyl substituted with 0-2 R³³,
 bicyclo[3.1.1]heptane substituted with 0-2 R³³,
 bicyclo[2.2.1]heptane substituted with 0-2 R³³, and
 phenyl substituted with 0-3 R³³;

- 10 R¹³ is H, methyl, or ethyl;

R³³, at each occurrence, is independently selected from

- H, F, Cl, Br, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl,
 t-butyl, -OH, methoxy, ethoxy, n-propoxy, i-propoxy, -SCH₃, -SCH₂CH₃,
 15 -SO₂CH₃, -CF₃, -OCF₃, CF₂CF₃, -CN, -NO₂, -NH₂, -CH₂NH(CH₃),
 -CH₂N(CH₃)₂, -NH(CH₃), -N(CH₃)₂, -CO(CH₃), -CO(OCH₃),
 -NHCO(CH₃), -CONH₂, -C(=O)H, -CH(OH)CH₃, -CH₂OH, -CH₂CH₂OH,
 -CH₂OCH₃, -CH₂CH₂OCH₃, and -CH₂OCH₂CH₃.

- 20 5. A compound of Claim 4 of Formula (Ib):



(Ib)

- 25 or a stereoisomer or a pharmaceutically acceptable salt form thereof, wherein:

R¹ is H or methyl;

R⁵ is H or methyl;

5 R⁶ is selected from

-F, -Cl, -CF₃, -CF₂CF₃, -OCF₃, -OCF₂CF₃, -OCF₂H, -OCF₂CH₃, -CN,
-OCH₃, -SCH₃, -S(=O)CH₃, -S(=O)₂CH₃, or methyl;

R⁷ is H, F, or Cl;

10

R⁸ is selected from

-OR¹², -SR¹², -NR¹²R¹³,
C₁₋₆ alkyl substituted with 0-2 R^{8a}, and
C₃₋₆ cycloalkyl substituted with 0-2 R^{8a},

15

R^{8a}, at each occurrence, is independently selected from

H, F, Cl, Br, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl,
t-butyl, -OH, methoxy, ethoxy, n-propoxy, i-propoxy, -CF₃, -OCF₃,
-CN, -CF₂CF₃, -SCH₃, -SCH₂CH₃, -CH₂NH(CH₃), -CH₂N(CH₃)₂,
20 -NH(CH₃), -N(CH₃)₂, -CO(CH₃), -CO(OCH₃), -NHCO(CH₃), -CONH₂,
-CH(OH)CH₃, -CH₂OH, -CH₂CH₂OH, -CH₂OCH₃, -CH₂CH₂OCH₃,
-CH₂OCH₂CH₃, phenyl substituted with 0-5 R³³, and
pyridyl substituted with 0-5 R³³

25 R⁹ is H;

R¹² is selected from

methyl, ethyl, propyl, butyl, pentyl, hexyl,
cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
30 bicyclo[3.1.1]heptane, bicyclo[2.2.1]heptane,

methyl substituted with R^{12a};
 ethyl substituted with R^{12a};
 propyl substituted with R^{12a};
 phenyl substituted with 0-2 R³³; and
 5 pyridyl substituted with 0-2 R³³;

R^{12a} is selected from


cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
 bicyclo[3.1.1]heptane, bicyclo[2.2.1]heptane, and
 10 phenyl substituted with 0-2 R³³;

R¹³ is H, methyl, or ethyl;

R³³, at each occurrence, is independently selected from

15 H, F, Cl, Br, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl,
 t-butyl, -OH, methoxy, ethoxy, n-propoxy, i-propoxy, -SCH₃, -SCH₂CH₃,
 -SO₂CH₃, -CF₃, -OCF₃, -CN, -NO₂, -NH₂, -CH₂NH(CH₃),
 -CH₂N(CH₃)₂, -NH(CH₃), -N(CH₃)₂, -CO(CH₃), -CO(OCH₃),
 -NHCO(CH₃), -CONH₂, -C(=O)H, -CH(OH)CH₃, -CH₂OH, -CH₂CH₂OH,
 20 -CH₂OCH₃, -CH₂CH₂OCH₃, and -CH₂OCH₂CH₃.

6. A compound of Claim 1 of Formula (Ic):



(Ic)

25

or a pharmaceutically acceptable salt thereof.

7. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of Claim 1 or
5 a pharmaceutically acceptable salt thereof.

8. A method for treating a human suffering from a disorder associated with 5HT_{2C} receptor modulation comprising administering to a patient in need thereof a therapeutically effective amount of a compound of Claim 1 or a pharmaceutically
10 acceptable salt thereof.

9. A method of Claim 8 for treating a human suffering from a disorder associated with 5HT_{2C} receptor modulation wherein the compound is a 5HT_{2C} agonist.

10. A method for treating obesity comprising administering to a patient in need thereof a therapeutically effective amount of a compound of Claim 1 or a
15 pharmaceutically acceptable salt thereof.